

References

1. Pomés A, Helm RM, Bannon GA, Burks AW, Tsay A, Chapman MD. Monitoring peanut allergen in food products by measuring Ara h 1. *J Allergy Clin Immunol* 2003;111:640-5.
2. Pomés A, Vinton R, Chapman MD. Peanut allergen (Ara h 1) detection in foods containing chocolate. *J Food Prot.* 2004 Apr; 67 (4):793-8.
3. Perry TT, Conover-Walker MK, Pomés A, Chapman MD, Wood RA. Distribution of peanut allergen in the environment. *J Allergy Clin Immunol.* 2004 May;113(5):973-6.
4. Maloney JM, Chapman MD, Sicherer SH. Peanut allergen exposure through saliva: assessment and interventions to reduce exposure. *J Allergy Clin Immunol.* 2006 Sep; 118(3):719-24.



www.inbio.com

Indoor Biotechnologies, Inc.
1216 Harris Street,
Charlottesville,
VA 22903
United States

Tel: (434) 984-2304
Fax: (434) 984-2709
E-mail: mail@inbio.com

Indoor Biotechnologies Ltd.
Cardiff Medicentre
Heath Park
Cardiff CF14 4UJ
Wales, United Kingdom

Tel: +44 (0) 2920 68 2115
Fax: +44 (0) 2920 68 2044
E-mail: info@indoorbiotech.co.uk




Ara h 6 ELISA kit (3B8/3E12)

Product Code: EL-AH6

Lot Number: xxxxx

Sample Curve:



Content:

- Vial 1 (red top) 100 μ L
Monoclonal antibody 3B8
Concentration: 2mg/ml in PBS
- Vial 2 (white top) 400 μ L
Ara h 6 Standard
Concentration: 1000ng/ml nAra h 6
- Vial 3 (brown) 100 μ L
Biotinylated monoclonal antibody 3E12
Dilute: 1:1000 for use

Storage: The ELISA kit should be stored at 4°C

For research and commercial use in vitro: not for human in vivo or therapeutic use.

Certificate of Analysis

Monoclonal Antibody: 3B8 (clone 3B8 B5)
Immunogen: Ara h 6
Isotype: Mouse IgG1
Specificity: Binds to species specific epitope present on *Arachis hypogaea* allergen, Ara h 6.
Purification: Produced in ascites and purified by chromatography using Protein G. Single heavy and light chain bands on SDS-PAGE.
Concentration: 2mg/ml in phosphate buffered saline, pH 7.4. Based on A280 for IgG (1.42=1mg/ml) 0.22µm filtered, preservative free.
Lot Number: xxxxx

Monoclonal Antibody: 3E12 (clone 3E12 C4 B3)
Immunogen: Ara h 6
Isotype: Mouse IgG1
Specificity: Binds to species specific epitope present on *Arachis hypogaea* allergen, Ara h 6.
Purification: Produced in ascites and purified by chromatography using Protein G. Single heavy and light chain bands on SDS-PAGE.
Biotinylation: Biotinylated and titrated for use in ELISA at 1/1000 dilution. Prepared in 1% BSA/50% glycerol/PBS, pH 7.4, 0.22µm filtered, preservative free.
Lot Number: xxxxx

Allergen Standard: nAra h 6
Composition: Naturally purified Ara h 6 prepared in 1% BSA, 50% glycerol/PBS, pH 7.4
Concentration: 1000 ng/ml nAra h 6
Calibration: The concentration of the purified natural Ara h 6 was determined by amino acid analysis.
Lot Number xxxxx

ELISA Protocol for Ara h 6

1. Coat polystyrene microtiter plates (NUNC Maxisorp Cert. NUNC catalog # 439454) with 100µl mAb 3B8 at 10µl/10ml, i.e. 1/1000 dilution of stock, in 50mM carbonate-bicarbonate buffer, pH 9.6, incubate overnight at 4°C.
2. Wash wells 3x with PBS-0.05% Tween 20, pH 7.4 (PBS-T). Incubate for 30 min. at room temperature with 100µl/well of 1% BSA, PBS-T. Wash 3x with PBS-T.
3. Use doubling dilutions of the nAra h 6 standard to make a control curve ranging from 100 - 0.2ng/ml Ara h 6: Pipette 20µl Ara h 6 standard into 180µl 1% BSA, PBS-T into wells A1 and B1 on the ELISA plate. Mix well and transfer 100µl across the plate into 100µl 1% BSA, PBS-T diluent to make 10 serial doubling dilutions. Wells A11, B11 and A12, B12 should contain only 1% BSA, PBS-T as blanks.
4. Add 100µl of diluted allergen samples and incubate for 1 hour at room temperature. House dust extracts for Ara h 6 analysis are routinely diluted two-fold from 1/10-1/80. Other sample types, like air filter extracts and allergen extracts, may require different dilutions.
5. Wash wells 3x with PBS-T and add 100µl diluted biotinylated anti Ara h 6 mAb 3E12. The antibody solution contains 50% glycerol and should be diluted 1/1000 in 1%BSA, PBS-T. Incubate for 1 hour at room temperature.
6. Wash wells 3x and add 100µl diluted Streptavidin - Peroxidase (Sigma S5512, 0.25mg reconstituted in 1ml distilled water). The Streptavidin Peroxidase should be diluted 1/1000 in 1% BSA, PBS-T. Incubate for 30 minutes at room temperature.
7. Wash wells 3x and develop the assays by adding 100µl 1mM ABTS in 70mM citrate phosphate buffer, pH 4.2 and 1/1000 dilution of H₂O₂. Read the plate when the absorbance at 405nm reaches 2.0-2.4.

Notes:

The Ara h 6 standard is recommended for immunoassay calibration purposes only. Not recommended for in-vitro antibody measurements, T cell studies, immunization purposes, or other uses.

Buffer recipes, storage conditions and a list of frequently asked questions can be found under "Protocols" on our web site: www.inbio.com.

For research and commercial use in vitro: not for human in vivo or therapeutic use.