

References:

1. Phipatanakul W, Eggleston PA, Wright EC, Wood RA. Mouse allergen. I. The prevalence of mouse allergen in inner-city homes. The National Cooperative Inner-City Asthma Study. *J Allergy Clin Immunol.* 2000;106:1070-4.
2. Phipatanakul W, Eggleston PA, Wright EC, Wood RA. Mouse allergen. II. The relationship of mouse allergen exposure to mouse sensitization and asthma morbidity in inner-city children with asthma. *J Allergy Clin Immunol.* 2000;106:1075-80.
3. Permaul P, Hoffman E, Fu C, Sheehan W, Baxi S, Gaffin J, Lane J, Bailey A, King E, Chapman M, Gold D, Phipatanakul W. Allergens in urban schools and homes of children with asthma. *Pediatr Allergy Immunol.* 2012;23:543-9.
4. Ahluwalia SK, Peng RD, Breyse PN, Diette GB, Curtin-Brosnan J, Aloe C, Matsui EC. Mouse allergen is the major allergen of public health relevance in Baltimore City. *J Allergy Clin Immunol.* 2013;132:830-5.
5. Ownby DR. Will the real inner-city allergen please stand up? *J Allergy Clin Immunol.* 2013;132:836-7.



www.inbio.com

Indoor Biotechnologies, Inc.
1216 Harris Street,
Charlottesville,
VA 22903
United States

Tel: (434) 984-2304
Fax: (434) 984-2709
E-mail: mail@inbio.com

Indoor Biotechnologies Ltd.
Cardiff Medicentre
Heath Park
Cardiff CF14 4UJ
Wales, United Kingdom

Tel: +44 (0) 2920 68 2115
Fax: +44 (0) 2920 68 2044
E-mail: info@indoorbiotech.co.uk




Mus m 1 ELISA kit

Product Code: EL-MM1

Lot Number: xxxxx

Sample Curve:



Contents:

- Vial 1 (red top) 100 μ L
Polyclonal rabbit anti-Mus m 1 capture antibody
Concentration: 1.6mg/ml in PBS
- Vial 2 (white top) 400 μ L
Universal Allergen Standard
Concentration: 250ng/ml Mus m 1
- Vial 3 (brown) 100 μ L
Biotinylated polyclonal rabbit anti-Mus m 1 antibody
Dilute: 1:1000 for use

Storage: The ELISA kit should be stored at 4°C

For research and commercial use in vitro: not for human in vivo or therapeutic use.

Certificate of Analysis

Antibody: Rabbit polyclonal antiserum
 Immunogen: Mus m 1
 Isotype: Multiple
 Specificity: Binds to an epitope on mouse *Mus musculus* urinary allergen, Mus m 1.
 Purification: Affinity chromatography using recombinant Protein G. Single heavy and light chain bands on SDS-PAGE
 Concentration: 1.6mg/ml in phosphate buffered saline, pH 7.4. Based on A280 for IgG (1.42=1mg/ml) 0.22µm filtered, preservative free.
Lot Number: xxxxx

Antibody: Rabbit polyclonal antiserum
 Immunogen: Mus m 1
 Isotype: Multiple
 Specificity: Binds to an epitope on mouse *Mus musculus* urinary allergen, Mus m 1.
 Purification: Affinity chromatography using recombinant Protein G. Single heavy and light chain bands on SDS-PAGE.
 Biotinylation: Biotinylated and titrated for use in ELISA at 1/1000 dilution. Prepared in 1% BSA/50% glycerol/PBS, pH 7.4, 0.22µm filtered, preservative free.
Lot Number: xxxxx

Allergen Standard: Universal Allergen Standard
 Composition: A formulation of eight purified natural allergens prepared in 1% BSA/50% glycerol/PBS, pH 7.4
Lot Number: xxxxx
 Concentration:

Universal Allergen Standard	Protein Measurement	Concentration (ng/ml)
Der p 1	Amino-acid analysis	2500
Der f 1	Amino-acid analysis	2500
Der p 2	Amino-acid analysis	1000
Fel d 1	Amino-acid analysis	1000
Can f 1	Amino-acid analysis	2500
Rat n 1	Amino-acid analysis	1000
Mus m 1	Amino-acid analysis	250
Bla g 2	Amino-acid analysis	2500

ELISA Protocol for Mus m 1

- Coat polystyrene microtiter plates (NUNC Maxisorp Cert. NUNC catalog # 439454) with 100µl pAb MM1 at 10µl/10ml, i.e. 1/1000 dilution of stock, in 50mM carbonate-bicarbonate buffer, pH 9.6, incubate overnight at 4°C.
- Wash wells 3x with PBS-0.05% Tween 20, pH 7.4 (PBS-T). Incubate for 30 min. at room temperature with 100µl/well of 1% BSA, PBS-T. Wash 3x with PBS-T.
- Use doubling dilutions of the Universal Allergen Standard to make a control curve ranging from 25 - 0.05ng/ml Mus m 1: Pipette 20µl Universal Allergen Standard into 180µl 1% BSA, PBS-T into wells A1 and B1 on the ELISA plate. Mix well and transfer 100µl across the plate into 100µl 1% BSA, PBS-T diluent to make 10 serial doubling dilutions. Wells A11, B11 and A12, B12 should contain only 1% BSA, PBS-T as blanks.
- Add 100µl of diluted allergen samples and incubate for 1 hour at room temperature. House dust extracts for Mus m 1 analysis are routinely diluted two-fold from 1/10-1/80. Other sample types, like air filter extracts and allergen extracts, may require different dilutions.
- Wash wells 3x with PBS-T and add 100µl diluted biotinylated pAb MM1. The antibody solution contains 50% glycerol and should be diluted 1/1000 in 1% BSA, PBS-T. Incubate for 1 hour at room temperature.
- Wash wells 3x and add 100µl diluted Streptavidin - Peroxidase (Sigma S5512, 0.25mg reconstituted in 1ml distilled water). The Streptavidin Peroxidase should be diluted 1/1000 in 1% BSA, PBS-T. Incubate for 30 minutes at room temperature.
- Wash wells 3x and develop the assays by adding 100µl 1mM ABTS in 70mM citrate phosphate buffer, pH 4.2 and 1/1000 dilution of H₂O₂. Read the plate when the absorbance at 405nm reaches 2.0-2.4.

Notes:

Universal Allergen Standard is recommended for immunoassay calibration purposes only. Not recommended for in-vitro antibody measurements, T cell studies, immunization purposes, or other uses.

Buffer recipes, storage conditions and a list of frequently asked questions can be found under "Protocols" on our web site: www.inbio.com.

For research and commercial use in vitro: not for human in vivo or therapeutic use.